



# Whole genome sequencing, clinical interpretation, and deep brain stimulation in a severely mentally ill person

Gholson J. Lyon<sup>1,2,3\*</sup>, Jason A. O'Rawe<sup>1,2</sup>, Han Fang<sup>1,2</sup>, Reid Robison<sup>3</sup>, Edward S. Kiruluta<sup>4</sup>, Gerald Higgins<sup>5</sup>, Martin G. Reese<sup>4</sup>

<sup>1</sup>Stanley Institute for Cognitive Genomics, One Bungtown Road, Cold Spring Harbor Laboratory, NY, USA; <sup>2</sup>Stony Brook University, 100 Nicolls Rd, Stony Brook, NY, USA; <sup>3</sup>Utah Foundation for Biomedical Research, E 3300 S, Salt Lake City, Salt Lake City, UT, USA; <sup>4</sup>Omicia Inc., 2200 Powell St., Emeryville, CA, USA; <sup>5</sup>AssureRx Health, Inc., 6030 S. Mason-Montgomery Road, Mason, Ohio, USA.

## Background

There is a rich literature in clinical psychology, psychiatry and neurology of single patient studies. Until now, however, such studies have focused on the parallel areas of clinical neuropsychiatry, personal genomics and brain-machine interfaces.

## Methods



Detailed phenotyping and clinical evaluations were conducted over a four-year period for a single United States Veteran male with severe mental illness. His genome was sequenced in the Illumina Whole Genome Sequencing Clinical Laboratory Improvement Amendments (CLIA)-certified laboratory. This person was implanted with the Medtronic Reclaim® Deep Brain Stimulation

(DBS) Therapy device for Obsessive Compulsive Disorder (OCD). Programming of the device and psychiatric assessments occurred in an outpatient setting for over two years.


## Results

We report here the detailed phenotypic characterization, clinical-grade whole genome sequencing (WGS), and two year outcome of a man with severe obsessive compulsive disorder treated with deep brain stimulation (DBS) targeting the nucleus accumbens / anterior limb of the internal capsule (ALIC). Since implantation, this man has reported steady improvement, highlighted by a steady decline in his Yale-Brown Obsessive Compulsive Scale (YBOCS) score from ~38 to a score of ~25. A rechargeable Activa RC neurostimulator battery has been of major benefit in terms of facilitating a degree of stability and control over the stimulation. His psychiatric symptoms reliably worsen within

hours of the battery becoming depleted, thus providing confirmatory evidence for the efficacy of DBS for OCD in this person. Whole genome sequencing revealed that he is a heterozygote for the p.Val66Met variant in BDNF, encoding a member of the nerve growth factor family, and which has been found to predispose carriers to various psychiatric illnesses. He carries the p.Glu429Ala allele in methylenetetrahydrofolate reductase (MTHFR) and the p.Asp7Asn allele in ChAT, encoding choline O-acetyltransferase, with both alleles having been shown to confer an elevated susceptibility to psychoses. We have found thousands of other variants in his genome, including pharmacogenetic variants, and have archived and offered the clinical sequencing data to him, so that he and others can re-analyze his genome for years to come. As this individual is a U.S. Veteran, we are working with the VA to incorporate his genomic data into the electronic medical record, VistA, which is of relevance to the One Million Veteran Program.



**Figure 1. Variant prioritization was performed on all variants discovered by the Illumina CLIA WGS pipeline using the Omicia Opal platform.** Variants were imported into the Omicia Opal cloud based clinical annotation and variant prioritization platform, and subsequently prioritized by requiring each variant to have prior evidence in OMIM and by additionally requiring each variant to be scored as having an Omicia Score of greater than 0.7.



**Figure 2. Yale Brown Obsessive Compulsive Scale (YBOCS) scores were measured for M.A. over a three year and seven months period of time.** A time series plot (A) shows a steady decline in YBOCS scores over the period of time spanning his DBS surgery (s) and treatment. Incremental adjustments to neurostimulator voltage are plotted over a period of time following DBS surgery (A). Mean YBOCS scores are plotted for sets of measurements taken before and after his Deep Brain Stimulation (DBS) surgery (B). A one-tailed unpaired t test with Welch's correction results in a p value of 0.0056, demonstrating a significant difference between YBOCS scores measured before and after the time of surgery.

## Conclusions

To our knowledge, this is the first study in the clinical neurosciences including 1) clinical-grade WGS with management and return of genetic results for a person with severe mental illness and 2) detailed neuropsychiatric phenotyping and individualized treatment with deep brain stimulation for his OCD. His WGS results and positive outcome with DBS for OCD is one example of individualized medicine in neuropsychiatry, including genomics-guided preventive efforts and brain-implantable devices. This is also an example of the split model for clinical genomics involving separate clinical-grade processes for sample collection, sequencing, analysis, and clinical interpretation. This serves as a model for the One Million Veteran Program, as this is the first genome sequenced and data returned for a U.S. Veteran, to our knowledge.

Questions? Email: [GholsonJLyon@gmail.com](mailto:GholsonJLyon@gmail.com)

Gene name	Genomic coordinates	Amino acid change	Zygosity	Mutation type	Population Frequency	Clinical significance
MTHFR	chr1:11854476	Glu>Ala	heterozygous	non-synon	T:77% G:23%	Susceptibility to psychoses, schizophrenia, occlusive vascular disease, neural tube defects, colon cancer, acute leukemia, and methylenetetra-hydrofolate reductase deficiency
BDNF	chr11:27679916	Val>Met	heterozygous	non-synon	C:77% T:23%	Susceptibility to OCD, psychosis, and diminished response to exposure therapy
CHAT	chr10:50824117	Asp>Asn	heterozygous	non-synon	G:85% A:15%	Susceptibility to schizophrenia and other psychopathological disorders.

**Table 1. A summary of three clinically relevant alleles found in the sequencing results of M.A.** Mutations in *MTHFR*, *BDNF*, and *CHAT* were found to be of potential clinical relevance for this person, as they are all implicated in contributing to the susceptibility and development of many neuropsychiatric disorders that resemble those present within M.A. A brief summary of the characteristics of each mutation is shown, including the gene name, genomic coordinates, amino acid change, zygosity, mutation type, estimated population frequency and putative clinical significance. Out of 91 other people in Utah genotyped on Illumina 2.5M arrays, only 2 of them carried all three of these variants, which is consistent with the noted minor allele frequencies on these 3 separate autosomes.